

## Crotoxin-Treated Macrophages Stimulate ROS Production and Killing Activity in co-cultured Neutrophils

Renata Begliomini Bruno De-Oliveira<sup>1#</sup>  
Ellen Emi Kato<sup>1#</sup>  
Maria Cristina Cirillo<sup>1#</sup>  
Tatiane Soares De-Lima<sup>1,2#</sup>  
Tatiana Carolina Alba-Loureiro<sup>3#</sup>  
Rui Curi<sup>4#</sup>  
Sandra Coccuzzo Sampaio<sup>1,5#\*</sup>

<sup>1</sup>Laboratory of Pathophysiology, Butantan Institute, Brazil

<sup>2</sup>Department of Molecular Biology and Biochemistry, Institute for Immunology, University of California Irvine, USA

<sup>3</sup>Department of Physiology and Biophysics, Institute of Biomedical Sciences, University of São Paulo, Brazil

<sup>4</sup>Interdisciplinary Post-Graduate Program in Health Sciences, Cruzeiro do Sul University, Brazil

<sup>5</sup>Department of Pharmacology, Institute of Biomedical Sciences, University of São Paulo, Brazil

### Abstract

Crotoxin (CTX), the predominant toxin in *Crotalus durissus terrificus* snake venom (CdtV), has anti-inflammatory and immunomodulatory effects. Despite its inhibitory action on neutrophil migration and phagocytosis, CTX does not directly affect the production of reactive oxygen species (ROS) by the neutrophils. In contrast, it enhances the generation of reactive oxygen and nitrogen intermediates by macrophages. Given the importance of macrophage-neutrophil interactions in innate antimicrobial defense, the aim of this study was to investigate the effect of CTX on neutrophil ROS production and killing activity, either through CTX-treated macrophage co-culture or conditioned medium of CTX-treated macrophages. The results showed an important modulatory action of CTX on the neutrophil function as well as neutrophil-macrophage interactions, as demonstrated by the increased production of hydrogen peroxide, hypochlorous acid, nitric oxide and TNF- $\alpha$ , along with the increased fungicidal activity of neutrophils.

**Keywords:** Crotoxin, Macrophages, Neutrophils, Inflammation, ATP, Reactive oxygen and nitrogen species, Cytokines, Co-culture model.

### Introduction

Macrophage-neutrophil interactions involve reciprocal activation of both cells [1,2] and play a key role in antimicrobial defense [2,3]. Neutrophils and macrophages have certain unique and complementary characteristics which allow them to cooperatively act as effectors and modulators of innate immunity [2,4]. The basic mechanism of macrophage-neutrophil co-operation is the transfer of microbicidal compounds such as ROS, RNS, and cytokines from the activated-macrophages to the neutrophils, which modulate the latter's response [1].

Neutrophils produce higher levels of ROS than macrophages. In addition, other antimicrobial factors like myeloperoxidase are either absent or present in scarce amounts in the monocytes, which results in a less intense microbicidal capacity compared to neutrophils[1,5,6]. During an inflammatory response against pathogens, co-operation between macrophages and neutrophils improves ROS production via the NADPH oxidase complex, and also increases the amounts of reactive nitrogen species (RNS), particularly nitric oxide (NO), leading

### Article Information

**Article Type:** Research

**Article Number:** JAMBR108

**Received Date:** 06 June, 2018

**Accepted Date:** 11 June, 2018

**Published Date:** 15 June, 2018

**\*Corresponding author:** Dr. Sandra Coccuzzo Sampaio, Department of Pharmacology, Institute of Biomedical Sciences, University of São Paulo, Av Prof Lineu Prestes, 1524, 05508-900, SP, Brazil. Tel: +55 11 2627-9562; Email: [sandra.coccuzzo\(at\)butantan.gov.br](mailto:sandra.coccuzzo(at)butantan.gov.br)

<sup>#</sup>These Authors are Joint First Authors on this Work.

**Citation:** De-Oliveira RBB, Kato EE, Cirillo MC, De-Lima TS, Alba-Loureiro TC, Curi R, Sampaio SC (2018) Crotoxin-Treated Macrophages Stimulate ROS Production and Killing Activity in co-cultured Neutrophils. J Appl Microb Res. Vol: 1, Issu: 1 (55-64).

**Copyright:** © 2018 Sampaio SC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

to the formation of more potent antimicrobial molecules like peroxy nitrite [3,7]. Macrophage-neutrophil interaction also involves the participation of PRRs (PAMPs receptors) that are expressed on both cells [8], and is mediated by cytokines such as IL-8, CCL2, IL-1  $\beta$  and TNF-  $\alpha$  [9,10] which further amplify the inflammatory response [11].

Several studies have characterized the anti-inflammatory and immunomodulatory potential of Crotoxin (CTX), the main toxic component of the *Crotalus durissus terrificus* snake venom [12-15]. First isolated by Slotta and Fraenkel-Conrat in 1938 [16], and structurally elucidated by Fraenkel-Conrat and Singer in 1956 [17], CTX is a  $\beta$ -neurotoxin heterodimer formed by non-covalent association of two different subunits—crotoxin (CA) and a phospholipase A<sub>2</sub> (PLA<sub>2</sub>-CB). Sixteen different isoforms of CTX were identified as a result of randomized combinations of the CA (CA1, CA2, CA3, and CA4) and CB (CBa2, CBb, CBc, and CBd) isoforms. The combinations of these isoforms form different complexes with different biological and pharmacological properties [18].

Both *in-vivo* and *in-vitro* studies have shown the dual effects of CTX on macrophage functions, such as inhibition of macrophage spreading and phagocytosis, and stimulation of H<sub>2</sub>O<sub>2</sub> and NO production and release [13,19]. These changes are accompanied by increased activity of the key glycolytic enzymes and higher consumption of glucose and glutamine, which demonstrates the metabolic effects of this toxin [20]. CTX also enhances the production of ROS, RNS, and cytokines in macrophages, thereby inducing the M1 or inflammatory phenotype, and enhancing their antitumor and anti-leishmanial properties [13,14]. In contrast, CTX inhibits both migration and phagocytosis of neutrophils induced by flogistic agents such as carrageenan or FMLP, and down-regulates the Syk-GTPase pathway [15,21]. This inhibitory effect of CTX on neutrophil functions contributes to its anti-inflammatory role that has been previously reported [12,15]. In addition, and in contrast to what has been observed for macrophages, CTX does not affect neutrophil metabolism (i.e. production of ROS induced by PMA) nor its microbicidal activities as seen in a model of peritonitis induced by carrageenan [21].

Since the CTX selectively stimulates macrophages, and the interaction between macrophages and neutrophils results in reciprocal stimulation, the question arises as to whether inflammatory macrophages treated with CTX are capable of altering the metabolism and function of co-cultured neutrophils. Therefore, the aim of the present study was to investigate whether the CTX induced macrophage activation induces a more potent fungicidal response by the neutrophils. We were able to demonstrate an enhanced *in-vitro* fungicidal activity of neutrophils against *Candida albicans* after co-culturing with CTX-treated macrophages. Our findings highlight the potential of this toxin in studying the mechanisms involved in the genesis of the inflammatory response, and in the regulation of neutrophils and macrophage interactions in innate immunity.

## Materials and Methods

### Animals

Male Wistar rats weighing 160-180 grams were used in this study. All animal experiments were performed in accordance with the guidelines for animal experimentation, and the Ethical Committee for the Use of Animals of the Butantan Institute approved the protocol (CEUAIB, protocol number 832/11 and 1013/13).

### Crotoxin

Crude venom solution was subjected to anion-exchange chromatography as previously described by Rangel-Santos *et al.* [22] using a Mono-Q HR 5/5 column in an FPLC system (Pharmacia, Uppsala, Sweden), and different fractions (1 mL/min) were eluted using a linear gradient of NaCl (0-1 mol/L in 50 mmol/L Tris-HCl, pH 7.0) [23]. Three peaks (p1, p2 and p3) were obtained of which p2 corresponded to the pure CTX fraction (about 60% of the crude venom), and peaks 1 and 3 included the other CdtV toxins. Prior to pooling, the fractions containing CTX were tested for homogeneity by non-reducing SDS-PAGE (12.5%) [24] and the phospholipase A<sub>2</sub> activity was assessed by a colorimetric assay using a synthetic chromogenic substrate, as previously described by our group [13,19,25,26].

### Peritoneal macrophages preparation

As per the recommendations of the Animal Ethics Committee Protocols (832/11 and 1013/13), the rats were euthanized in a CO<sub>2</sub> chamber and their peritoneal cavity was opened and washed with 10 mL cold phosphate-buffered saline (PBS) pH 7.4. After gently massaging the abdominal wall, the peritoneal fluid now containing the resident macrophages was aspirated. The total number of peritoneal cells was counted and differential counts were performed on smears stained with a panchromatic dye [27]. The samples from individual animals were used for all the measurements. The assays were always performed in duplicate as described in our previous studies [13,19,25,26].

### Peritoneal neutrophilic exudate

Peritoneal exudates were obtained as described previously by our group [21]. The rats were injected intraperitoneally with carrageenan (Cg-4.5 mg/kg) (Sigma, St Louis, MO, USA) and euthanized four hours later. The peritoneal cavity was washed with PBS as described, and after a gentle massage of the abdominal wall, the peritoneal fluid containing neutrophils was aspirated. The total and differential peritoneal cell counts [27] indicated that 95% of the cells were neutrophils [21].

### Treatment with CTX and neutrophil-macrophage co-cultures

The neutrophil-macrophage co-culture protocol was adapted from the method described by Hauptmann *et al.* [28]. Resident peritoneal macrophages were suspended in RPMI 1640 medium supplemented with 10% fetal calf serum and antibiotics (penicillin 50 U/ml and streptomycin 50  $\mu$ g/mL) at the density of  $4 \times 10^6$ /mL and seeded into a 24-well plate. After letting the cells to adhere for an hour, they were

washed and incubated with CTX (0.3  $\mu\text{g/mL}$  or 12.5 nM) for 2 h under 5%  $\text{CO}_2$  at 37 °C. The cells were washed once to remove the CTX and cultured for 24 h with fresh medium containing 5 $\mu\text{g/mL}$  LPS, and the LPS conditioned medium (CM) was collected. Peritoneal inflammatory neutrophils were suspended at the density of  $4 \times 10^6/\text{mL}$  in either the CM (Neutr  $\phi$  :CM-Macr  $\phi$  contact) for monocultures, in complete RPMI medium and seeded onto the inflammatory macrophage monolayers for co-cultures (Neutr  $\phi$  :Macr  $\phi$  contact), or in complete medium with 5  $\mu\text{g/mL}$  LPS for control monocultures. After 24 h of culture, the neutrophils were harvested to measure ATP,  $\text{H}_2\text{O}_2$ , HOCl, NO and cytokine levels and fungicidal activity. All experiments were performed in triplicate with macrophages and neutrophils from three different animals. The concentration of CTX and the time of incubation with the toxin were the same as those used in previous studies [13,19,21,25,26], and were not cytotoxic as per Trypan blue exclusion test Figure 1.

### Hydrogen peroxide production

The production of  $\text{H}_2\text{O}_2$  was measured as described by Pick *et al.* [29], based on a horseradish peroxidase (HRP) dependent conversion of phenol red into a colored compound by  $\text{H}_2\text{O}_2$ . After 24 h of culture, the harvested neutrophils were re-suspended at the density of  $4 \times 10^6$  cells/mL in phenol red solution (PRS) containing 140 mM NaCl, 10 mM potassium-phosphate buffer pH 7.0, 5 mM dextrose, 0.28 mM phenol red, and 8.5 U/mL HRP to a final volume of 7.4 mL with Hank's solution. PRS solution was used as blanks, serial dilutions of  $\text{H}_2\text{O}_2$  from 5 to 40  $\mu\text{M}$  were prepared in PRS for the standard curves, and neutrophil suspension with 100 ng phorbol myristate acetate (PMA) was used as the positive control. One hundred microliters of each solution were plated per well of a 96-well flat-bottomed tissue culture plate (Corning, NY®), and incubated in a humidified atmosphere at 37 °C for 1 h. The reaction was stopped by the addition of 10  $\mu\text{L}$  of 1 N NaOH. The  $\text{H}_2\text{O}_2$  dependent phenol red oxidation was spectrophotometrically measured at 620 nm using a

Titertek Multiscan apparatus. The concentration of  $\text{H}_2\text{O}_2$  was expressed as nano-moles of  $\text{H}_2\text{O}_2$  per  $4 \times 10^5$  cells.

### Hypochlorous acid production

Hypochlorous acid production (HOCl) was measured as described by Dypbukt *et al.* [30]. After 24 h of co-culture, neutrophils were harvested and  $4 \times 10^5$  cells were re-suspended in 250  $\mu\text{L}$  of 5 mM taurine (Sigma) solution containing 140 mM NaCl, 10 mM KCl, 0.5mM  $\text{MgCl}_2$ , 1 mM  $\text{CaCl}_2$ , 1 mg/mL glucose and 100 ng/mL PMA, and incubated for 1 h at 37°C under 5%  $\text{CO}_2$ . To evaluate the basal production of HOCl, a PMA-minus control was included. The reaction was stopped by the addition of 20  $\mu\text{g/mL}$  catalase (Sigma), and the samples were immersed in an ice bath for 10 min. After centrifuging the samples, the supernatants were collected and 50  $\mu\text{L}$  of 2 mM 3,3',5,5'-tetramethylbenzidine (TMB; Sigma) solution containing 10  $\mu\text{M}$  KI and 10% DMF in 400 mM acetate buffer pH 5.4 was added to each sample. After five minutes of incubation, the reduction of TMB was measured spectrophotometrically at 650 nm, against a blank solution consisting of 5 mM taurine solution. HOCl concentration was measured with the help of a standard curve and expressed as *micro*-moles of HOCl per  $4 \times 10^5$  neutrophils per hour.

### Determination of extracellular ATP concentration

Harvested neutrophils were re-suspended in 100  $\mu\text{L}$  1% Triton-X, and lysed by heat shock (alternating exposure to hot and cold temperatures three times) to release ATP. After centrifuging the lysates, 10  $\mu\text{L}$  of the supernatants were pipetted into each well of a flat white-bottomed 96 well plate (Corning, Costar). Serial dilutions of the ATP solution provided by the manufacturer were prepared for the standard curve. The ATP Determination Kit (A22066, Molecular Probes®) was used according to the manufacturer's instructions. Briefly, 0.1 M DTT (dithiothreitol) was added to each well followed by 100  $\mu\text{L}$  of the 10  $\mu\text{M}$  luciferase/luciferin reagent solution. The release in ATP was measured for 10 min at 20 s intervals, at the maximum emission of 560 nm against the




Figure 1: Scheme of the experimental design.

reaction buffer. ATP concentration was calculated with the help of the standard curve, and expressed as nano-moles of ATP per  $1 \times 10^6$  cells, as described by Sakaki *et al.* [31].

### Nitric oxide production

For measuring NO production, the harvested neutrophils were re-suspended in fresh medium with or without *C. albicans*, and incubated for 4 h. Nitrite levels in the supernatants were measured as described by Ding *et al.* [32]. Briefly, 50  $\mu$  L of each supernatant was incubated with an equal volume of Griess reagent (1% sulfanilamide, 0.1% naphthalene diamine dihydrochloride, 2.5%  $H_3PO_4$ ) at room temperature for 10 min. The absorbance at 550 nm was measured against the cell-free medium containing 0.2-0.3  $\mu$  M  $NO_2^-$  as blank. The nitrite concentration was determined with the help of sodium nitrite standard curves.

### Cytokine quantification

To determine cytokine production, the harvested neutrophils were re-suspended in fresh medium with or without *C. albicans* and incubated for 4 h. Cytokines in the supernatants were quantified using ELISA as per the manufacturer's instructions. Briefly, ELISA plates (Immuno Maxisorp; Nunc<sup>®</sup>, NJ) were coated with mouse anti-rat monoclonal antibodies (anti-TNF- $\alpha$  IgG1 Clone # 45418 and anti-CINC-2 IgG2b Clone # 123802) or polyclonal antibody (anti-IL-1 $\beta$ ; R&D Systems, Minneapolis, MN), and incubated overnight at room temperature. After blocking for 1 h at room temperature, the respective samples and standards were added and incubated for 2 h. Biotinylated secondary antibodies were then added and incubated for another 2 h followed by a 20 min incubation with peroxidase-conjugated streptavidin. The color was developed using TMB in the dark for 20 min. The reaction was stopped with 2N sulfuric acid, and absorbance was measured at 465 and 590 nm in a microplate reader (Spectra Max 190, Molecular Devices). The cytokine concentrations were determined with the help of the standard curve prepared using recombinant murine cytokines (R&D Systems), with a detection sensitivity of 4-10 pg/mL.

### Phagocytosis of *C. albicans* and fungicidal activity

*C. albicans* (ATCC-9002-8) was cultured in 10% Sabouraud's dextrose broth (Microbiology and Mycology Laboratories, Department of Clinical Analyses, Faculty of Pharmaceutics Science, University of São Paulo) at 30 °C for 24 h. The fungi were centrifuged, washed twice with Dulbecco PBS, and re-suspended at  $2 \times 10^7$ /mL. The viability was determined by methylene blue 0.05% (>98%) exclusion, and the cells counted in a Neubauer's chamber. Phagocytosis and fungicidal activity were measured as described by Sampaio *et al.* [33] and Lima *et al.* [21]. Neutrophils harvested from the co-cultures or control monocultures ( $1 \times 10^6$  cells/mL) were incubated in sterile plastic tubes (to avoid cell adherence) at 37 °C with *C. albicans* ( $4 \times 10^6$  fungi/mL) in RPMI 1640 medium supplemented with 10% FCS. After 40 min of incubation, 200  $\mu$  L of the neutrophil-yeast suspension were adhered onto glass coverslips by cytocentrifugation. The coverslips were stained with Wright's and May-Giemsa,

and the fungicidal activity was determined by assessing cell viability of phagocytosed particles by the dye exclusion test [27]. The number of candidacidal neutrophils was scored as 0 (no *Candida* cells killed), 1 (1 or 2 cells), 2 (3 or 4 cells), and 3 (>4 cells). The index of candidacidal activity was calculated by the sum of the scores obtained per animal, as described by Sampaio *et al.* [33].

### Statistical analysis

All statistical analyses of the differences between the groups were performed according to Glantz [34] using GraphPad InStat software, version 3.01 (GraphPad Software Inc., San Diego, CA, USA). A one-way analysis of variance followed by Tukey's test was used for multiple comparisons (all pairs of groups) of values from the assays comparing the monoculture and co-culture groups. To analyze data of the other assays, ANOVA followed by Bonferroni's test was used for multiple comparisons against a single control, and unpaired Student's *t*-test was used to compare two groups. P values < 0.05 were considered statistically significant. The results were presented as the mean values  $\pm$  standard error.

## Results

### Crotoxin-treated macrophages stimulate hydrogen peroxide release in co-cultured neutrophils


To determine the effect of CTX on ROS production, neutrophils, and macrophage mono- and co-cultures were treated with the toxin, and  $H_2O_2$  levels were measured. Neutrophils co-cultured with LPS and CTX-stimulated macrophages (Neutr  $\phi$  :Macr  $\phi$  contact) or cultured with the conditioned medium collected from these macrophages (Neutr  $\phi$  :CM-Macr  $\phi$  contact) showed increased levels of  $H_2O_2$  compared to neutrophils co-cultured with untreated macrophages (Figures 2A and 2B). In contrast, while CTX increased  $H_2O_2$  production in macrophages (Figure 2C), it did not stimulate  $H_2O_2$  production by neutrophil monocultures under the same conditions (Figure 2D), thus corroborating our previous data [21].

### Crotoxin-treated macrophages stimulate hypochlorous acid production in co-cultured neutrophils


Neutrophils co-cultured with LPS and CTX stimulated macrophages showing the significant increase in basal HOCl production (38%), as well as that after PMA stimulation (76%) (Figure 3A). Similar effects were observed in neutrophils cultured with the CM from LPS and CTX stimulated macrophages (Figure 3B). However, CTX alone did not stimulate HOCl production by neutrophils (Figure 3C).

### Crotoxin-treated macrophages stimulate ATP production in co-cultured neutrophils


Neutrophils co-cultured with LPS and CTX treated macrophages or the CM of stimulated macrophages showed increased production of ATP by 52% and 42%, respectively (Figure 4). Macrophages treated with the toxin significantly increased ATP production by 91% whereas CTX alone did not affect ATP production in neutrophils (Figure 4).



**Figure 2:** Release of hydrogen peroxide by neutrophils co-cultured with CTX-treated macrophages. The release of H<sub>2</sub>O<sub>2</sub> was measured in neutrophils ( $4 \times 10^6$  cells) co-cultured for 24 h with LPS-stimulated macrophages ( $4 \times 10^6$  cells/LPS- $5\mu\text{g/ml}$ ) that were pretreated with CTX ( $12.5 \text{ nM}$ ) for 2 h (Neutr  $\phi$  :Macr  $\phi$ ) (A) or in the macrophage-conditioned supernatant (Neutr  $\phi$  :CM-Macr  $\phi$ ) (B). In panels (C) and (D), the release of H<sub>2</sub>O<sub>2</sub> was evaluated in monocultures of CTX-treated macrophages (Macr  $\phi$  monoculture) and neutrophils (Neutr  $\phi$  monoculture), respectively, maintained under the same experimental conditions. The assays were performed in quadruplicates and samples without PMA were evaluated to assess the spontaneous production of H<sub>2</sub>O<sub>2</sub>. The absorbance was determined in ELISA reader, with a filter of 620 nm obtaining the optical density (O.D.) and calculating the results in nmoles of H<sub>2</sub>O<sub>2</sub> per  $4 \times 10^5$  cells. The results express the mean  $\pm$  e.p.m of three animals for each cell type and represent three different experiments. \*P<0.001.



**Figure 3:** Release of hypochlorous acid by neutrophils co-cultured with CTX-treated macrophages. The release of H<sub>2</sub>O<sub>2</sub> was measured in neutrophils ( $4 \times 10^6$  cells) co-cultured for 24 h with LPS-stimulated macrophages ( $4 \times 10^6$  cells/LPS- $5\mu\text{g/ml}$ ) that were pretreated with CTX ( $12.5 \text{ nM}$ ) for 2 h (Neutr  $\phi$  :Macr  $\phi$ ) (A) or in the macrophage-conditioned supernatant (Neutr  $\phi$  :CM-Macr  $\phi$ ) (B). In panel (C), the release of HOCl was evaluated in monocultures of CTX-treated neutrophils (Neutr  $\phi$  monoculture), maintained under the same experimental conditions. The assays were performed in quadruplicates and samples without PMA were evaluated to assess the spontaneous production of HOCl. The absorbance was determined in ELISA reader, with a filter of 650 nm obtaining the optical density (O.D.) and calculating the results in  $\mu\text{M}$  HOCl per  $4 \times 10^5$  cells. The results express the mean  $\pm$  e.p.m of three animals for each cell type and represent three different experiments. \*P<0.001.



**Figure 4:** Release of ATP by neutrophils co-cultured with CTX-treated macrophages. Neutrophils (number) were co-cultured for 24 h with LPS-stimulated macrophages that were pretreated with CTX for 2 h (number, LPS dose, CTX dose) or in the macrophage-conditioned supernatant. Monocultures of CTX-treated macrophages and neutrophils were also analyzed, maintained under the same experimental conditions. The release of ATP was measured by an integration time of 10 min (20 s intervals) at a maximum emission of 560 nm and the results were expressed by a standard curve of extracellular ATP levels, in nmoles of ATP/1 × 10<sup>6</sup> cells. The results express the mean ± e.p.m of three animals for each cell type and represent two different experiments. \*P<0.05, compared to the value of the respective control group.

### CTX-treated macrophages increased neutrophil killing activity by inducing an early release of TNF- $\alpha$ and nitric oxide


Neutrophils co-cultured with LPS and CTX stimulated macrophages or CM increased their fungicidal capacity by 30% and 42%, respectively, compared to CTX-untreated co-cultures. However, CTX did not augment the fungicidal activity of neutrophil monocultures (Figure 5). To determine the underlying mechanism of the higher fungicidal action induced by CTX, we evaluated the cytokine secretion and NO production in the suitable cultured neutrophils after a 4 h incubation with *C. albicans*.

The neutrophils co-cultured with the stimulated macrophages or the CM showed increased NO production by 103% and 101%, respectively, while the CTX treated neutrophil monocultures did not show any change in NO production after 4 h of fungal exposure (Figure 6A). Interestingly, there was no change in the secreted cytokine profile of neutrophils after 24 h of incubation in the CM-Macrophage stimulated with CTX and LPS (*data not shown*). However, since cytokines are modulated at the beginning of neutrophil-macrophage interaction during antimicrobial defense (Ref), we determined whether a shorter co-culture period between neutrophils and macrophages resulted in a distinct cytokine secretion profile during phagocytosis of *C. albicans*. Neutrophils co-cultured for 4 h with stimulated macrophages showed no changes in the levels of IL-1  $\beta$  and CINC-2 (Figure 6B and 6C), but secreted 4.25 higher levels of TNF- $\alpha$  during the candidacidal period compared to neutrophils co-cultured with unstimulated macrophages (Figure 6D).


### Discussion

Based on the functional complementarity between macrophages and neutrophils, and on the distinct effects of CTX on the metabolism of these cells, the modulatory action of macrophages treated with CTX on ROS production and fungicidal activity of neutrophils was evaluated. Our results demonstrated that inflammatory neutrophils co-cultured with LPS and CTX stimulated macrophages, or with the supernatant from these macrophages had increased metabolism and fungicidal capacity. In addition, CTX also intensified the metabolism of LPS-stimulated macrophages. Our data are consistent to the findings of Faiad *et al.* [35] who demonstrated that CTX increased ROS and RNS production in both resident and tumor macrophages obtained from the peritoneal cavity of tumor-bearing rats. Therefore, we demonstrated that the stimulatory action of CTX on macrophage metabolism is independent of the stimulus involved.


Several studies have demonstrated that LPS activates M1 macrophages to respond to injury [36], and are characterized by increased production and release of reactive oxygen and nitrogen species, compounds that are known to enhance neutrophil secretory capacity [37-39]. In fact, the co-culture of neutrophils with LPS-stimulated macrophages increased the latter's inflammatory activity. In a previous study, our group demonstrated that CTX could not directly modulate neutrophil metabolism [21]. However, the peculiar metabolic state of macrophages induced by CTX was able to alter neutrophil metabolism, and stimulate these cells to produce more ROS and ATP, and the latter in turn is known to increase the production of multiple inflammatory mediators like superoxide anion, hydrogen peroxide, nitric



**Figure 5:** Fungicidal capacity of neutrophils co-cultured with CTX-treated macrophages. This activity was determined in neutrophils (number) obtained from monocultures or after co-cultured with LPS-stimulated macrophages that were pretreated with CTX for 2 h (number, LPS dose, CTX dose). The assay was determined by an exclusion staining test, in which only live intracellular yeasts are stained in blue. Live and dead yeasts were counted in a total of 100 cells and the results were expressed as a score. The results express the mean  $\pm$  e.p.m of three animals for each cell type and represent three different experiments. \*P<0.05.



**Figure 6:** Cytokine production by neutrophils during fungicidal activity. Neutrophils (number) were co-cultured with LPS-stimulated macrophages pretreated with CTX (number, LPS dose, CTX dose) or in conditioned medium from these macrophages. In panel (A), NO production from Neutrφ monoculture treated or not with CTX and Neutrφ co-cultured with Macrφ or in Mφ-CM pretreated with CTX. In panel (B, C, and D), cytokine production by Neutrφ monoculture treated or not with CTX, maintained under the same experimental conditions and by Neutrφ co-cultured with Macrφ or in Mφ-CM pretreated with CTX. The assays were performed in triplicate. In the NO production assay, the absorbance was determined by an ELISA reader, at 540 nm. The results are expressed in  $\mu$ moles of NO<sub>2</sub> per  $4 \times 10^5$  cells. In the cytokine production assay, the absorbance was determined in an ELISA reader at 450 nm and calculating the results in pg/mL. The results are expressed as the mean  $\pm$  e.p.m of three animals for each cell type and represent three different experiments. \*P<0.001.



oxide, and cytokines, among others [40-42]. Therefore, the increase in ATP is an important parameter of neutrophil metabolism as well as activity. Our study has demonstrated for the first time the ability of CTX to induce ATP production in macrophages as well as in neutrophils co-cultured with the stimulated macrophages, without affecting the viability of either. This is consistent with the study of Faiad *et al.* [20] who demonstrated the stimulatory action of CTX on glucose and glutamine oxidation in rat peritoneal macrophages, which led to an increase in ATP production [43]. The findings of both studies reinforce the stimulatory capacity of CTX on the metabolism of macrophages and the importance of ATP in neutrophil activation during the interaction between these cells. Currently, our group is studying the mechanisms involved in CTX mediated activation of macrophages and neutrophils.

The activation of neutrophils culminates in the production of NO and pro-inflammatory cytokines such as IL-1 β, IL-8, and TNF- α, which further amplifies the interaction between neutrophils and macrophages during inflammatory responses against infections [1,44]. The fungicidal activity of neutrophils may involve a more modulated cytokine response. For example, TNF- α is quickly produced in response to *C. albicans* infection [45], which is known to activate macrophages and neutrophils by increasing their phagocytic capacity and production of ROS and RNS, which eliminate the invading pathogens [46]. Therefore, we evaluated the production of cytokines and NO by neutrophils cultured with supernatants obtained from the CTX-treated macrophages as a measure of their fungicidal activity. The neutrophils showed a significant increase in TNF- α levels, which is consistent with previous data that show the appearance of this cytokine in the early stages of neutrophil-macrophage interaction during the inflammatory response against infections [47]. Interestingly, the NO production was significantly increased in the co-cultured neutrophils

during the fungicidal phase. Therefore, we hypothesize that the increase in TNF- α production may have contributed to the increase in NO production, which in turn enhanced the fungicidal ability of neutrophils. Peroxynitrite (ONOO<sup>-</sup>), a product of spontaneous NO and O<sub>2</sub> fusion, is one of the most important oxidizing agents in various pathophysiological conditions, and could also be generated in neutrophils during CTX-treated macrophage co-culture, leading to increased fungicidal ability [48-51].

Our group has also recently demonstrated that CTX-treatment induced an M1 profile in macrophages infected with *Leishmania*, with increased ROS, NO, IL-6 and TNF- α production and without loss of viability [14], which is associated with a better prognosis for cutaneous leishmaniasis treatment. Lipoxins (LXA) and their lipid derivatives are potent stimulators of the microbicidal activity of macrophages and increase the levels of intracellular ROS [52-56]. Recent studies on both *in vitro* and *in vivo* models have shown a stimulatory effect of CTX on the production of LXA<sub>4</sub> and its stable analog 15-Epi-LXA<sub>4</sub>, along with increased ROS and RNS production, in resident macrophages or those are the part of the tumor microenvironment [13,25,57]. Therefore, we hypothesize that these mediators secreted by CTX-treated macrophages, regardless of their activation state, may enhance the fungicidal capacity of co-cultured neutrophils. In fact, secretion of LXA<sub>4</sub> and 15-Epi-LXA<sub>4</sub>, and anti-inflammatory mediators like resolvins and protectins, by macrophages in an infectious microenvironment regulate the inflammatory response by inhibiting neutrophil recruitment [58,59]. Therefore, if we consider the ability of CTX to stimulate both pro- and anti-inflammatory mediators, it is likely that the activation of neutrophils by CTX-treated macrophages would increase the fungicidal response without exacerbating local inflammation. In fact, the modulatory capacity of CTX can be attributed to its pleiotropic effects on different inflammatory mediators [13].

Our study demonstrates for the first time that increased ROS production and fungicidal activity of neutrophils are associated with increased oxidative metabolism in CTX-treated macrophages. LPS-stimulated macrophages induce production of ROS, ATP, RNS, and TNF- α by neutrophils during the early stages of contact (H<sub>2</sub>O<sub>2</sub> release from 1 hour of incubation - Figure 1 of the *Supplementary Material*), and the high levels are maintained for an extended duration (Figure 7). This could be relevant to the modulation of neutrophils during the infectious inflammatory response *in-vivo*.

## Conclusion

Therefore, this study contributes to the knowledge on CTX modulatory action on the inflammatory response and highlights once again the potential of this toxin for studying the mechanisms involved in inflammatory response vis-a-vis neutrophil and macrophage interactions in innate immunity.

## Acknowledgment

This work was supported by the grants 2009/52330-9, São Paulo Research Foundation (FAPESP) and 2008/57898-0, Instituto Nacional de Ciência e Tecnologia em Toxinas (INCTTOX) and Guggenheim Foundation; fellowships

2012/08880-7, São Paulo Research Foundation (FAPESP) and Secretaria da Saúde do Estado de São Paulo (PAP-SSP). The authors would like to thank Mr. Andre Fonseca Alves for his valuable technical assistance with the purification of CTX.

**Conflict of interest:** The authors declare that there are no conflicts of interest.

## References

- Silva MT (2010) Neutrophils and macrophages work in concert as inducers and effectors of adaptive immunity against extracellular and intracellular microbial pathogens. *J Leukoc Biol* 87: 805-813.
- Silva MT (2010) When two is better than one: macrophages and neutrophils work in concert in innate immunity as complementary and cooperative partners of a myeloid phagocyte system. *J Leukoc Biol* 87: 93-106.
- Silva MT (2011) Macrophage phagocytosis of neutrophils at inflammatory/infectious foci: a cooperative mechanism in the control of infection and infectious inflammation. *J Leukoc Biol* 89: 675-683.
- Machado RL, Araujo SA, Carvalho L, Carvalho EM (2004) Immune response mechanisms to infections. *Am bras Dermatol* 79: 647-664.
- Levy O (2004) Antimicrobial proteins and peptides: anti-infective molecules of mammalian leukocytes. *J Leukoc Biol* 76: 909-925.
- Segal AW (2005) How neutrophils kill microbes. *Annu Rev Immunol* 23: 197-223.
- Tharmalingam S, Alhasawi A, Appanna VP, Lemire J, Appanna VD (2017) Reactive nitrogen species (RNS)-resistant microbes: adaptation and medical implications. *Biol Chem* 398: 1193-1208.
- Akira S, Takeda K, Kaisho T (2001) Toll-like receptors: critical proteins linking innate and acquired immunity. *Nat Immunol* 2: 675-680.
- Cassatella MA, Gasperini S, Calzetti F, McDonald PP, Trinchieri G (1995) Lipopolysaccharide-induced interleukin-8 gene expression in human granulocytes: transcriptional inhibition by interferon-gamma. *Biochem J* 310: 751-755.
- Kasama T, Miwa Y, Isozaki T, Odai T, Adachi M, et al. (2005) Neutrophil-derived cytokines: potential therapeutic targets in inflammation. *Curr Drug Targets Inflamm Allergy* 4: 273-279.
- Scapini P, Lapinet-Vera JA, Gasperini S, Calzetti F, Bazzoni F, et al. (2000) The neutrophil as a cellular source of chemokines. *Immunol Rev* 177: 195-203.
- Nunes FP, Zychar BC, Della-Casa MS, Sampaio SC, Gonçalves LR, et al. (2010) Crotoxin is responsible for the long-lasting anti-inflammatory effect of *Crotalus durissus terrificus* snake venom: involvement of formyl peptide receptors. *Toxicon* 55: 1100-1106.
- Costa ES, Faiad OJ, Landgraf RG, Ferreira AK, Brigatte P, Curi R, et al. (2013) Involvement of formyl peptide receptors in the stimulatory effect of crotoxin on macrophages co-cultivated with tumour cells. *Toxicon* 74: 167-178.
- Farias LHS, Rodrigues APD, Coelho EC, Santos MF, Sampaio SC, et al. (2017) Crotoxin stimulates an M1 activation profile in murine macrophages during *Leishmania amazonensis* infection. *Parasitology* 144: 1458-1467.
- Lima TS, Neves CL, Zambelli VO, Lopes FSR, Sampaio SC, et al. (2017) Crotoxin, a rattlesnake toxin, down-modulates functions of bone marrow neutrophils and impairs the Syk-GTPase pathway. *Toxicon* 136: 44-55.
- S KH, HF-C (1938) Estudos químicos sobre venenos ofídicos. Entregue para publicação.
- Fraenkel-Conrat H, Singer B (1956) Fractionation and composition of crotoxin. *Arch Biochem Biophys* 60: 64-73.
- Faure G, Xu H, Saul FA (2011) Crystal structure of crotoxin reveals key residues involved in the stability and toxicity of this potent heterodimeric  $\beta$ -neurotoxin. *J Mol Biol* 412: 176-191.
- Sampaio SC, Brigatte P, Sousa-e-Silva MC, dos-Santos EC, Rangel-Santos AC, et al. (2003) Contribution of crotoxin for the inhibitory effect of *Crotalus durissus terrificus* snake venom on macrophage function. *Toxicon* 41: 899-907.
- OJ F (2011) Crotoxin stimulates glucose metabolism in peritoneal macrophages during tumor progression.
- Lima TS, Cataneo SC, Iritus AC, Sampaio SC, Della-Casa MS, et al. (2012) Crotoxin, a rattlesnake toxin, induces a long-lasting inhibitory effect on phagocytosis by neutrophils. *Exp Biol Med* 237: 1219-1230.
- Rangel-Santos A, Dos-Santos EC, Lopes-Ferreira M, Lima C, Cardoso DF, et al. (2004) A comparative study of biological activities of crotoxin and CB fraction of venoms from *Crotalus durissus terrificus*, *Crotalus durissus cascavella* and *Crotalus durissus collilineatus*. *Toxicon* 43: 801-810.
- Lôbo de Araújo A, Radvanyi F (1987) Determination of phospholipase A2 activity by a colorimetric assay using a pH indicator. *Toxicon* 25: 1181-1188.
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
- Sampaio SC, Alba-Loureiro TC, Brigatte P, Landgraf RG, Dos Santos EC, et al. (2006) Lipoxygenase-derived eicosanoids are involved in the inhibitory effect of *Crotalus durissus terrificus* venom or crotoxin on rat macrophage phagocytosis. *Toxicon* 47: 313-321.
- Sampaio SC, Santos MF, Costa EP, Rangel-Santos AC, Carneiro SM, et al. (2006) Crotoxin induces actin reorganization and inhibits tyrosine phosphorylation and activity of small GTPases in rat macrophages. *Toxicon* 47: 909-919.
- G Rosenfeld (1971) *Symptomatology, Pathology and Treatment of snakes bites in South America*. Academic Press, New York.
- Hauptmann S, Zwadlo-Klarwasser G, Jansen M, Klosterhalfen B, Kirkpatrick CJ (1993) Macrophages and multicellular tumor spheroids in co-culture: a three-dimensional model to study tumor-host interactions. Evidence for macrophage-mediated tumor cell proliferation and migration. *Am J Pathol* 143: 1406-1415.
- Pick E, Mizel D (1981) Rapid micro assays for the measurement of superoxide and hydrogen peroxide production by macrophages in culture using an automatic enzyme immunoassay reader. *J Immunol Methods* 46: 211-226.
- Dypbukt JM, Bishop C, Brooks WM, Thong B, Eriksson H (2005) A sensitive and selective assay for chloramine production by myeloperoxidase. *Free Radic Biol Med* 39: 1468-1477.
- Sakaki H, Tsukimoto M, Harada H, Moriyama Y, Kojima S (2013) Autocrine regulation of macrophage activation via exocytosis of ATP and activation of P2Y11 receptor. *PLoS One* 8: e59778.
- Ding AH, Nathan CF, Stuehr DJ (1988) Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages. Comparison of activating cytokines and evidence for independent production. *J Immunol* 141: 2407-2412.
- Sampaio SC, Sousa-e-Silva MC, Borelli P, Curi R, Cury Y (2001) *Crotalus durissus terrificus* snake venom regulates macrophage metabolism and function. *J Leukoc Biol* 70: 551-558.
- Stanton G (1997) *Primer of Bio-Statistics*. McGraw-Hill, New York.
- Faiad OJ (2012) Efeito da Crotoxina sobre função e o metabolismo de glicose e glutamina de macrófagos durante a progressão tumoral.
- Al-Shabany AJ, Moody AJ, Foey AD, Billington RA (2016) Intracellular NAD<sup>+</sup> levels are associated with LPS-induced TNF- $\alpha$  release in pro-inflammatory macrophages. *Biosci Rep* 36: e00301.
- Marcinkiewicz J, Czajkowska B, Grabowska A, Kasprovicz A, Kociszewska B (1994) Differential effects of chlorination of bacteria on their capacity to generate NO, TNF-alpha and IL-6 in macrophages. *Immunology* 83: 611-616.
- Sierra-Filardi E, Puig-Kröger A, Blanco FJ, Nieto C, Bragado R, et al. (2011) Activin A skews macrophage polarization by promoting a proinflammatory phenotype and inhibiting the acquisition of anti-inflammatory macrophage markers. *Blood* 117: 5092-5101.

39. Muralidharan S, Mandrekar P (2013) Cellular stress response and innate immune signaling: integrating pathways in host defense and inflammation. *J Leukoc Biol* 94: 1167-1184.
40. Gordon S (1986) Biology of the macrophage. *J Cell Sci Suppl* 4: 267-286.
41. Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci USA* 97: 8841-8848.
42. Verkhatsky A, Burnstock G (2014) Biology of purinergic signaling: its ancient evolutionary roots, its omnipresence and its multiple functional significance. *Bioessays* 36: 697-705.
43. Spolarics Z, Lang CH, Bagby GJ, Spitzer JJ (1991) Glutamine and fatty acid oxidation are the main sources of energy for Kupffer and endothelial cells. *Am J Physiol* 261: G185-90.
44. Kobayashi Y (2010) The regulatory role of nitric oxide in pro-inflammatory cytokine expression during the induction and resolution of inflammation. *J Leukoc Biol* 88: 1157-1162.
45. Vecchiarelli A, Puliti M, Torosantucci A, Cassone A, Bistoni F (1991) *In vitro* production of tumor necrosis factor by murine splenic macrophages stimulated with mannoprotein constituents of *Candida albicans* cell wall. *Cell Immunol* 134: 65-76.
46. Sredni-Kenigsbuch D, Kambayashi T, Strassmann G (2000) Neutrophils augment the release of TNF $\alpha$  from LPS-stimulated macrophages via hydrogen peroxide. *Immunol Lett* 71: 97-102.
47. Møller K, Strauss GI, Qvist J, Fonsmark L, Knudsen GM, et al. (2002) Cerebral blood flow and oxidative metabolism during human endotoxemia. *J Cereb Blood Flow Metab* 22: 1262-1270.
48. Huie RE, Padmaja S (1993) The reaction of NO with superoxide. *Free Radic Res Commun* 18: 195-199.
49. Szabó C, Ischiropoulos H, Radi R (2007) Peroxynitrite: biochemistry, pathophysiology and development of therapeutics. *Nat Rev Drug Discov* 6: 662-680.
50. Pacher P, Beckman JS, Liaudet L (2007) Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87: 315-424.
51. Carballal S, Bartsaghi S, Radi R (2014) Kinetic and mechanistic considerations to assess the biological fate of peroxynitrite. *Biochim Biophys Acta* 1840: 768-780.
52. Schwab JM, Chiang N, Arita M, Serhan CN (2007) Resolving E1 and protecting D1 activate inflammation-resolution programmes. *Nature* 447: 869-874.
53. Serhan CN, Chiang N, Van Dyke TE (2008) Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol* 8: 349-361.
54. Serhan CN, Yang R, Martinod K, Kasuga K, Pillai PS, et al. (2009) Maresins: novel macrophage mediators with potent anti-inflammatory activity. *J Biol Chem* 284: 15700-15710.
55. matory and pro-resolving actions. *J Exp Med* 206: 15-23.
56. Spite M, Norling LV, Summers L, Yang R, Cooper D, et al. (2009) Resolving D2 is a potent regulator of leukocytes and controls microbial sepsis. *Nature* 461: 1287-1291.
57. Cash JL, Christian AR, Greaves DR (2010) Chemerin peptides promote phagocytosis in a ChemR23- and Syk-dependent manner. *J Immunol* 184: 5315-5324.
58. Brigatte P, Faiad OJ, Nocelli RCF, Landgraf RG, Palma MS, et al. (2016) Walker 256 Tumor Growth Suppression by Crotoxin Involves Formyl Peptide Receptors and Lipoxin A<sub>4</sub>. *Mediators of Inflammation* Article ID 2457532, 11 pages.
59. Kumar V, Sharma A (2010) Neutrophils: Cinderella of innate immune system. *Int Immunopharmacol* 10: 1325-1334.