

Association between Vitamin D Deficiency and Cardiovascular Risk Factors in Elderly

Catarina Magalhães Porto^{1*}Vanessa de Lima Silva²Brivaldo Markman Filho³Rita de Cássia Hoffmann Leão⁴Camila da Costa Lima Souto⁵Vera Magalhães⁶Francisco Bandeira⁷¹Department of Cardiology and Gerontology at Federal University of Pernambuco, Brazil²Department of Speech-Language Pathology and Audiology at Federal University of Pernambuco, Brazil³Department of Cardiology at Federal University of Pernambuco, Brazil⁴Department of Geriatrics at Federal University of Pernambuco, Brazil⁵Department of the Gerontology at Federal University of Pernambuco, Brazil⁶Department of the Infectious Diseases at Federal University of Pernambuco, Brazil⁷Department of the Endocrinology and Medical Sciences at University of Pernambuco (UPE), Brazil

Abstract

It is important to identify earlier cardiovascular risk factors in the elderly due to cardiovascular aging and associated chronic diseases. In addition to traditional cardiovascular risk factors, vitamin D deficiency is also a new risk factor. To evaluate the association between vitamin D deficiency with cardiovascular risk factors in elderly patients in outpatient cardiology clinics was performed Cross-sectional study. Clinical data were collected from patients aged over 60 years in cardiology clinics of UFPE, from November 2015 to March 2016. The variables were independent, vitamin D deficiency and dependent on age, sex, education, ethnicity, hypertension, diabetes mellitus, hypothyroidism, renal failure, dementia, stroke, dyslipidemia, depression, smoking, alcoholism, obesity, andropause, coronary artery disease, left ventricular hypertrophy or impaired left ventricular function and cardiac arrhythmia. They were used the chi-square test Person and, when necessary, Fisher's exact test. Among the 137 elderly, was identified 91.2% were hypertensive; 35.0% with definite or possible coronary artery disease, 27.7% with cardiac arrhythmia or left ventricular hypertrophy. 65% of the elderly were deficient in vitamin D. More vitamin D deficiency risk were male gender, age \leq 70 years, smokers, obese, diabetic, with scores compatible with dementia, depression, cardiac arrhythmia, coronary heart disease, ventricular hypertrophy left or reduction in left ventricular function. There was a high prevalence of vitamin D deficiency in the elderly, strongly associated with smoking, obesity, diabetes mellitus, dementia, depression, cardiac arrhythmia, coronary artery disease and left ventricular hypertrophy or impaired left ventricular function.

Keywords: Vitamin D, Aged, Risk.

Introduction

In addition to the cardiovascular risk factors common to individuals, in general, such as biological, lifestyle, metabolic and socioenvironmental, as regards the elderly population, are also the gradual reduction of cardiovascular reserves and the decline of functions endocrine diseases, increasing the risk of cardiovascular diseases [1].

Article Information

Article Type: Research**Article Number:** JCDM109**Received Date:** 31 May, 2018**Accepted Date:** 14 June, 2018**Published Date:** 26 June, 2018

***Corresponding author:** Dr. Catarina Magalhães Porto, Cardiologist at the Hospital das Clínicas of the Federal University of Pernambuco, Doctor of the Health Service of the Federal Public Ministry, Rua Jacobina, 45, Apto 2801, Graças, Recife-PE, Brazil. Tel: + (81) 32313135/96557408; Email: [catarinamqo\(at\)gmail.com](mailto:catarinamqo(at)gmail.com)

Citation: Porto CM, Silva VL, Filho BM, Leão RCH, Souto CCL, et al (2018) Association between Vitamin D Deficiency and Cardiovascular Risk Factors in Elderly. J Cardiovasc Dis Med. Vol: 1, Issu: 1 (43-52).

Copyright: © 2018 Porto CM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Due to the modifications of the cardiovascular system inherent to aging, as well as the chronic diseases associated with this system, the adoption of a healthy lifestyle, with earlier identification and reduction of cardiovascular risk factors, was encouraged [2].

The literature is extensive, reporting an association between vitamin D deficiency with a higher prevalence of cardiovascular endpoints such as angina, acute myocardial infarction (AMI), heart failure (HF) and peripheral vascular disease, compared to individuals with higher levels of vitamin D. In addition, large prospective studies have also demonstrated this association between calcitriol deficiency and increased cardiovascular risk [3-7], as well as associated with higher risk of death [6,8,9] and a comprehensive systematic review and meta-analysis [10] due to the probable anti-inflammatory and antiatherogenic mechanisms of this vitamin, suggesting that vitamin D deficiency is a cardiovascular risk factor [11].

The 2010 Consensus for Vitamin D Nutrition Guidelines Consensus noted that more than 50% of older people worldwide do not have satisfactory levels of vitamin D [12].

High concentrations of vitamin D3 predict thirteen fewer years of mortality and cardiovascular disease (CVD) incidence [13]. A recent July 2015 study evaluating the longevity of centenarians and their relationship with vitamin D levels, comparing with patients with early AMI and healthy individuals from the same age group, found that centenarians had higher levels of vitamin D than normal adults and concluding that these levels of vitamin D because they confer physiological advantages and better cardiovascular function, are associated with successful aging [14].

This study has as an object the association between vitamin D deficiency and sociodemographic and cardiovascular risk factors in the elderly attended at a cardiology outpatient clinic.

Materials and Methods

It is an Epidemiological Study, Cross-Sectional and Analytical. The study sites were the Center of Attention to the Elderly of the Federal University of Pernambuco (UFPE) and the cardiology outpatient clinic of the UFPE Hospital das Clinicals. The selection of the population of this study was carried out from the analysis of records of the elderly, attended in these services, from November 2015 to March 2016.

The adopted age cut obeyed the Brazilian legislation, regardless of the health status of the person, according to Law No. 8.842 of 1994 [15]. Assuming a prevalence of 80% of vitamin D deficiency among the elderly, a drawing effect equal to one, the sample size equaled 137 records, corresponding to a 95% test power.

For inclusion in the study, it was admitted that they should be medical records of elderly patients, who presented for routine cardiac evaluation, attended by the researcher in a cardiology outpatient clinic, by referral or spontaneous demand, regardless of sex and motor restrictions.

The exclusion criterion was absence of record of all parameters of interest in the medical record or record of the elderly presenting a state of health compromised with indication of hospitalization, at the cardiological examination.

The study variables are characterized according to Table 1.

The instruments of data collection were the copy of the protocol of cardiological investigation used in the ambulatory of Cardiology of the places of study, plus the Mental State Mini-Exam and the Geriatric Depression Scale. After approval of the research project by the Ethics Committee for Research on Human Beings, data collection consisted of requesting the File and Statistical Service of the study sites the patient records that obeyed the inclusion criteria of this study.

A database with all the dependent variables and the independent variable of this research, recorded in the medical records, using the Epi7 program, version 7.1.5.2, of the WHO was elaborated. The data were analyzed with the IBM/SPSS Statistical Package for Social Sciences, version 20.0.

Hypothyroidism, renal insufficiency, dementia, cerebrovascular accident, cardiac arrhythmia, coronary artery disease (CAD), left ventricular hypertrophy, or reduction of left ventricular ejection fraction (LVEF) were considered as dependent variables: gender, education, marital status, ethnicity, depression, alcoholism, obesity, andropause, as well as the serum concentration of 25 hydroxyvitamin D was admitted as an independent variable measured by the chemiluminescence method (Architect Abbott®).

All variables on nominal or ordinal scales were summarized by absolute and relative frequency distributions, with subsequent dichotomization in order to integrate the inferential analyzes. In the bivariate analysis, Pearson's Chi-square test and, when necessary, the Fisher's exact test were used to test the association between the variables. All conclusions were taken using the significance level of 5%.

Ethical Considerations

This research obeyed the determinations of Resolution N° 466, of 2012, of the National Health Council, which governs research involving human beings and was approved by the Ethics Committee of the Federal University of Pernambuco under CAAE n° 47317715.6.0000.5208.

Data collection was authorized through the letters of consent from the Center for Elderly Care and Management of the Hospital das Clinicals of the Federal University of Pernambuco. The researcher undertook to maintain the confidentiality of the data and the identification of the patients in any publication.

Results

Analyzing the characteristics of the 137 elderly people surveyed, according to the data presented in Table 1, we found that 75.9% of them are female, 50.4% had brown skin

color, 62% are married or have a stable union, and 69.3 % belong to class C. Of the elderly, 66.4% have a minimum level of education up to full elementary education II. The mean age of the studied group was 71 years (standard deviation = 0.6), about 47% of them, almost half, were less than 70 years old (Table 2).

Analyzing the clinical variables of the present study, 78.8% of the elderly were overweight or obese, 72.3% had a waist/hip circumference index higher than normal and 94, 2% have dyslipidemia (Table 3).

Regarding the assessment of mental state, in the cognitive evaluation, 22.6% of the elderly had a mini-mental test compatible with dementia, and 31.4% had a moderate or severe depression. In the cardiovascular evaluation, practically the majority have hypertension (91.2%), 35% with CAD defined or possible and 27.7% with cardiac arrhythmia, and the same percentage has LVH (Table 3).

In order to verify the existence of association of the vitamin D concentration with the variables related to the socio-demographic characteristics of the elderly and the other clinical variables, Fisher's exact test was applied and presented in Table 4 the results in which this association

was significant. The elderly groups with the highest odds of presenting vitamin D deficiency were male, 70 years of age or older, smoking, obese, diabetic, with dementia score, with depression, cardiac arrhythmia, CAD and with LVH or Reduction of left ventricular ejection fraction

Discussion

The majority of the elderly in this study were overweight or obese, as well as having a waist index above normal, a high prevalence of dyslipidemia and systemic arterial hypertension, according to the literature, where aging is associated with a higher prevalence of alterations in glycemetic metabolism, central obesity, dyslipidemia and hypertension, characterizing Metabolic Syndrome [16].

Although vitamin D deficiency is well documented in elderly people living in countries of the northern hemisphere, in Brazil there is evidence of a high prevalence of vitamin D deficiency, as in a temperate climate study in São Paulo that demonstrated hypovitaminosis D in 43.5% of the active elderly, 87.6% of the elderly living in communities and 91.5% of those institutionalized [17]. In the present study conducted in Recife, northeast of the country, where the tropical climate predominates most of the time, we found a high percentage (65%) of vitamin D deficiency similar to

Table 1: Independent and dependent variables of the study, Hospital das Clínicas, Recife, 2015-2016.

Variables	Definition		Categorization
Independent variable	Vitamin D deficiency	< 30 ng/mL	Present/absent
Dependent variable	Sex	Informed by the patient, corresponding to last grade attended.	Male or female
	Schooling		Illiterate, Basic, middle, higher
	Marital Status		Single, married, Stable union, Widow/er, separated/
	Ethnicity	According to orientation from the Brazilian Institute of Geography and Statistics, based on self-referred color of skin.	White, black, Mixed colored, yellow, Native American, and dycotomized as white and non-white.
	IncomeClass Brasil	Evaluated based on possession of appliances, educational level of head of families and access to public services.	A, B1, B2, C1, C2, D, E
	Hypothyroidism	TSH> 4.7 mU/L (ages 60 and 69 years); 5.6 mU /L (ages 70 and 79 years); 6.30 mU /L (ages ≥ 80 years) – independent of the tetraiodothyronine hormone concentration and of the presence of suggestive clinical signs.	Present Absent
	Renal insufficiency	Glomerular filtration rate less than 60ml/min/1.73m ² associated to at least one marker of parenchymal renal damage present (Albuminuria > 30 mg / 24 hor glomerular hematuria)	Present Absent
	Dementia	Characterized by a score lower than 17 for illiterate patients, lower than 23 for patients with one to 11 years of education, and less than 28 when having more than 11 years of education.	Present Absent
	Stroke	Medical records of history and previous treatment of stroke.	Present Absent
	Depression	Mood alterations marked by irritability for most of the day, with slowing of mental functions and motor skills, and reduced ability of attention and concentration.	Present Absent
	Alcoholism	Score higher than 2 in the CAGE questionnaire.	Present Absent
	Obesity	Body mass index higher than 29.99 kg/m ² .	Present Absent
	Waist-hip ratio	Considered normal when less than 0.99 for men and 0.97 for women.	Present Absent
	Andropause	Total testosterone concentration lesser than 300 ng/dL associated to compatible clinical signs.	Present Absent
	Cardiac arrhythmia	Identified by the cardiac auscultation and resting lectrocardiogram and medical record entries.	Present Absent

Table 2: Distribution of sociodemographic characteristics of the elderly-NAI, Hospital das Clínicas, Recife, 2015-2016.

Sociodemographic characteristics	Categories	Frequency	Percentage
Sex	Male	33	24.1
	Female	104	75.9
Ethnicity	White	50	36.5
	Mixed color	69	50.4
	Black	18	13.1
Marital status	Single	10	7.3
	Married or stable union	85	62
	Widow/er	37	27
	Divorced or Separated	5	3.6
Education of the elder	Illiterate	11	8
	Basic I	39	28.5
	Basic II	41	29.9
	High school	30	21.9
	Higher education	16	11.7
Age classes (years)	60 - 69	64	46.7
	70 -79	60	43.8
	80 +	13	9.5
Economic class, Brazil	A	2	1.5
	B1	2	1.5
	B2	14	10.2
	C1	34	24.8
	C2	61	44.5
	D e E	24	17.5

Table 3: Frequency distribution of risk factors for cardiovascular disease and previous diseases of the elderly - NAI, Hospital das Clínicas, Recife, 2015-2016.

Risk factors and previous diseases	Categories	Frequency	Percentage
Smoking	Never	89	65
	Past	44	32.1
	Current	4	2.9
Risk of alcoholism (CAGE)	Yes	4	2.9
	No	133	97.1
Hormonal and metabolic measures			
Obesity	Normal weight	29	21.2
	Overweight	66	48.2
Waist-hip ratio	Obese	42	30.6
	Normal	16	11.7
Dyslipidemia	Absent	121	88.3
	Present	129	94.2
Andropause*	Absent	8	5.8
	Present	24	17.7
Diabetes mellitus	Absent	9	6.6
	Present	52	38
Hypothyroidism	Absent	85	62
	Present	8	5.8
Mental state			
Compatible score with dementia	Absent	129	94.2
	Present	31	22.6
Depression	Normal	106	77.4
	Moderate	94	68.6
	Grave	41	29.9
		2	1.5
Diagnosed conditions			
Systemical Hypertension SH	Absent	125	91.2
	Present	12	8.8
Chronic renal failure	Absent	4	2.9
	Present	133	97.1
Previous stroke	Absent	9	6.6
	Present	128	93.4

Cardiac Arrhythmia	Present	38	27.7
	Absent	99	72.3
Left ventricle Hypertrophy or reduction of the fraction of left ventricular ejection	Present	38	27.7
	Absent	99	72.3
Coronary artery disease	Absent	89	65
	Possible/Defined	48	35

Table 4: Association of sociodemographic and clinical variables with the concentration of Vitamin D – NAI, Hospital das Clínicas, Recife – 2016.

Variables	Category	Vitamin D concentration				Bivariate analysis	
		Deficient		Sufficient		OR (IC 95%)	p-valor
		N	%	N	%		
Sex	Male	28	84,8	5	15,2	3,948 (1,411-11,041)	0,006
	Female	61	58,7	43	41,3		
Age classes (years)	≤ 70	53	72,6	20	27,4	2,061 (1,010 - 4,205)	0,045
	> 70	36	56,3	28	43,8		
Ethnicity	White	35	70,0	15	30,0	1,426 (0,678 - 3,000)	0,349
	Not white	54	62,1	33	37,9		
Marital status	Other	36	69,2	16	30,8	1,358 (0,652 - 2,831)	0,413
	Married or stable union	53	62,4	32	37,6		
Education of the elder	Illiterate/ basic	64	70,3	27	29,7	1,991 (0,956 - 4,149)	0,064
	High school and Higher Education	25	54,3	21	45,7		
smoking	Current/Past	41	85,4	7	14,6	5,003 (2,027 - 12,348)	0,000
	Never	48	53,9	41	46,1		
Obesity	Obese	38	90,5	4	9,5	8,196 (2,711 - 24,780)	0,000
	Normal/Overweight	51	53,7	44	46,3		
Waist-hip ratio	Normal	62	62,6	37	37,4	1,465 (0,651 - 3,295)	0,355
	Increased	27	71,1	11	28,9		
Hypothyroidism	Present	8	100,0	0	0,0		0,051*
	Absent	81	62,8	48	37,2		
Chronic renal failure*	Present	4	100,0	0	0,0		0,297*
	Absent	85	63,9	48	36,1		
Previous stroke	Present	8	88,9	1	11,1	4,642 (0,563 - 38,276)	0,120
	Absent	81	63,3	47	36,7		
Diabetes	Present	42	80,8	10	19,2	3,396 (1,508 - 7,644)	0,002
	Absent	47	55,3	38	44,7		
Compatible score with dementia	Present	27	87,1	4	12,9	4,790 (1,565 - 14,665)	0,003
	Absent	62	58,5	44	41,5		
Depression	Moderate/severe	34	79,1	9	20,9	2,679 (1,155 - 6,214)	0,019
	Normal	55	58,5	39	41,5		
alcoholism*	Present	3	75,0	1	25,0	1,640 (0,166 - 16,205)	1,000*
	Absent	86	64,7	47	35,3		
Cardiac Arrhythmia	Present	30	78,9	8	21,1	2,542 (1,058 - 6,112)	0,034
	Absent	59	59,6	40	40,4		
Coronary artery disease	Possible/Defined	45	93,8	3	6,3	15,341 (4,438 - 53,032)	0,000
	No	44	49,4	45	50,6		
Left ventricle Hypertrophy or reduction of the fraction of left ventricular ejection	Present	37	97,4	1	2,6	33,442 (4,414 - 253,379)	0,000
	Absent	52	52,5	47	47,5		

NHANES III data [18].

The prevalence of hypovitaminosis D in the present study was also lower than that reported in the French population, mainly above 60 years (80.3%) [19], as well as in northern India, near the Himalaya, where the prevalence of vitamin D deficiency was 83%, associated with chronic stable angina considering the greater latitude of these countries [20], where the winters are longer and the individuals use clothes that cover practically the whole body, in contrast to the regions near the line of the Equator, Where sunny weather predominates most of the year.

Although there is no uniformity in the literature regarding cut-off points and dosage method, vitamin D deficiency is considered by the Institute of Medicine at a concentration of 25-hydroxyvitamin D less than 20 ng/mL and insufficiency when there are levels between 21 and 29 ng/ml [21].

However, this nomenclature changes. Thus, most studies consider deficient 25 (OH) D levels below 30 ng/mL, such as the Endocrine Society, the National Osteoporosis Foundation (NOF), the International Osteoporosis Foundation (IOF), and the American Geriatric Society (AGS). Severe deficiency when less than 20 ng/mL, but also the studies with results expressed in nmol/L classified the severe deficiency in the

presence of levels below 25 nmol/L, moderate of 25 to 49.9 nmol/L, insufficiency of 51 At 75 nmol/L [21,22]. In this present study, the presence of 25 (OH) D levels below 30ng/mL was considered as vitamin D deficiency.

As a justification for the high prevalence of vitamin D deficiency in the elderly, as related in this research, the literature exposes several factors, such as obesity, chronic liver or kidney diseases, diseases that impair fat absorption, lifestyle and environmental factors, such as institutionalization, reduction of outdoor physical activity, functional limitation [23], leading to less sun exposure and less outdoor activities, type of clothing, inadequate diet, use of medications that may interfere with vitamin D metabolism and, especially, reduced intake epidermal production of the vitamin D precursor due to skin atrophy due to aging [11].

The analysis of the previous clinical characteristics of the elderly sample of the present study underscores the importance of this approach, since about 30% of this elderly had a history of cardiac arrhythmia, Left ventricular hypertrophy (LVH) and CAD. In this study, a high prevalence of vitamin D deficiency was found in men, and it can be justified considering that women use vitamin D supplementation, because they are more affected than men by osteoporosis, as well as frequent health service are more subject to preventive medicine than men [24].

One of the limitations of this study did not quantify the elderly who were already using supplementation with vitamin D and it is worth mentioning that old women accompanied the services of this study are also accompanied by gynecologists and geriatricians, which are usually evaluated in a preventive or therapeutic as osteoporosis.

The clinical conditions presented by the elderly in this study, such as smoking, obesity, DM II, scores compatible with dementia, depression, cardiac arrhythmia, CAD, and LVH were related to a higher prevalence of vitamin D deficiency. The association between smoking and vitamin D deficiency, found in this study, may be related to inflammatory mechanisms present in chronic obstructive pulmonary disease (COPD) [25], as well as the functional limitations and greater sedentary lifestyle that these patients with COPD present, with less time of sun exposure, as well as by accelerated skin aging due to smoking, renal dysfunction and use of corticosteroids in COPD [26].

The association between obesity and vitamin D deficiency found in this study was also demonstrated in in another study, when 10.331 participants were cross-checked, finding an inverse association between waist circumference and calcidiol levels and systolic blood pressure [27]. It is worth mentioning that obese elderly people, due to their greater difficulty in locomotion, are less exposed to the sun, have less vitamin D₃ hepatic synthesis and are more absorbed by body fat deposits and thus have lower serum levels of 25 hydroxyvitamin D [28].

Regarding the association between vitamin D deficiency and diabetes in the present study, this was widely evaluated by several authors [29-32] as well as showing that the risk

of DM II can be reduced by 41% if levels of 25 (OH) D are sufficient [33].

The finding of the association between vitamin D deficiency and cognitive decline, assessed by the Mental State Minisex of this study was similar to that reported by Llewellyn et al. [34-37] with vitamin D being considered an essential neurohormonal hormone in the function of brain regulation and protection, against cognitive decline and Alzheimer's disease [38].

Depression is considered a cardiovascular risk factor, increased twice the risk of AMI and up to 2.5 years of cardiovascular morbidity and mortality [39] was correlated in this vitamin D-deficient research, being in concordance with works in the literature that found association between levels of vitamin D and depression [40-43]. A scientific research demonstrated the actions of this vitamin in the central nervous system, as well as the meta-analysis demonstrated a positive association between depression and vitamin D deficiency, indicates that vitamin supplementation may promote a significant improvement of this disorder and more recently, as demonstrated in a recent study conducted in teachers in Malaysia, where there was a high prevalence of vitamin D deficiency with low levels of this vitamin significantly associated with depression and worse quality of life [44-46].

The association between hypovitaminosis D and cardiac arrhythmia, mainly atrial fibrillation (AF) whose data were confirmed in the present study were also described by [6,47-50], and several mechanisms to explain this association are described, such as the negative regulation of calcitriol over Renin Angiotensin Aldosterone System (RAAS), reducing the oxidative stress, decreasing reactive oxygen species and inflammation [51], and prolonging the action potential through its role in the regulation of calcium metabolism [52]. However, other studies [53-55], based on the prospective cohort of the Rotterdam Study, evaluated the elderly with a 12-year follow-up, also did not find this association.

Although significant advances in primary prevention and treatment strategies for CAD, especially in the geriatric population, which are exposed to higher risks of worse outcomes after acute coronary syndrome, this remains the leading cause of death in developed countries [56].

In a recent study, evaluated octogenarians undergoing percutaneous coronary intervention, found that they had a 10.6 times higher risk of mortality after AMI with ST segment elevation, as well as acute stent thrombosis, previous AMI, Cardiac insufficiency, low left ventricular ejection fraction, ventricular arrhythmias and multiarterial lesions are the independent risk factors for in-hospital mortality in this population [57]. As explanation for these worse results of CAD in the elderly could be due to poor nutrition and low levels of vitamin D in this population favoring the pathogenesis of these complications [56].

The CAD in association with the vitamin D deficiency found in this study was related in many studies, such as the large prospective study of HPFS [3] and others such

as the Framingham Offspring Prospective Study [58], Cardiovascular Health Study [59], Augsburg Case-Cohort Study [5], which revealed an inverse association between vitamin D levels and CAD. The interpretation for the association between CAD and vitamin D deficiency may be due to possible inflammatory mechanism induced by Vitamin D deficiency leads to oxidative stress and release of proinflammatory cytokines causing platelet activation and release by the bone marrow of immature platelets with increased volume and activated with a higher rate of collagen aggregation, high levels of thromboxane A2, favoring greater expression of glycoproteins Ib and IIb/IIIa, leading to an increased risk of CAD [60].

It was observed that evaluating epicardial coronary flow in 222 patients submitted to coronary angiography due to suspicion of ischemic heart disease, thickening of the intima and averages carotid ultrasound as a marker of subclinical atherosclerosis and dilation of the brachial artery mediated flow as an indicator of endothelial dysfunction, found strong association between vitamin D insufficiency and coronary slow flow as well as endothelial dysfunction and subclinical atherosclerosis were more prevalent among subjects with vitamin D insufficiency [7].

As to the percentage of elderly patients with poor left ventricular systolic function or LVH associated with vitamin D deficiency, in the present study, this correlation with literature data was demonstrated [61], as well as related to pathophysiological mechanisms by which vitamin D acts in LVH through of VDR receptor, inhibiting RAAS [62,63]. Calcitriol, through its VDR receptor regulates the proliferation, morphology and growth of cardiomyocytes, and reduces the expression of the Atrial Natriuretic Factor and increases the expression and nuclear localization of the VDR in these cells [64,65].

Vitamin D exerts its anti-inflammatory role in LVH, reducing IL-2, TNF alpha, interferon gamma, TH1 lymphocytes [66] and reducing the expression of TNF-alpha and NF-kappa beta messenger RNA (In the present study, it was found that the TIMPs were able to inhibit the degradation of the extracellular matrix, which could prevent the evolution to HF [67]. VDR receptor type BSMT, which were related to LVH [68].

Was performed a cross-sectional study in elderly patients attending cardiology outpatient clinics of the UFPE, in Recife, from August 2015 to February 2016, in order to analyze the association between vitamin D deficiency and heart failure risk. Using bivariate logistic analysis, followed by multivariate logistic regression analysis, the authors found that among 137 elderly, the risk of heart failure was significantly associated with vitamin D deficiency, male gender, obesity and cardiac arrhythmia. The increased risk of heart failure in this study was present in more than half of the elderly and was significantly associated with vitamin D deficiency (growing 12.2 times the risk of heart failure) [69].

Cardiovascular diseases in the elderly have been considered a challenge for public health, being causes of high mortality in this population, despite the greater disclosure

of healthy aging, even in developed countries. Hence the importance of developing research to identify possible causes of cardiovascular risk, which could be directed to public health policies and actions and among such research, vitamin D deficiency.

The reduction of calcitriol leads to an increase in the production of PTH, extracellular matrix metalloproteinases (MMPs), Tumor Necrosis Factor alpha (TNF alpha), Advanced glycation end products (AGEs), reduction of carboxyglutamic acid matrix protein (matrix GLA protein) which is a dependent vitamin K, associated with the reduction of arterial calcification, reduction of osteopontin, of Interleukin-10 and type IV collagen, changes that lead to the formation of arterial calcification which in turn increases the risk of DCV [70]. The other mechanism of increased CVD-induced vitamin D deficiency is the increase in ANP (Natriuretic Atrial Peptide), changes in cardiomyocyte contractility, and decreased synthesis of extracellular matrix inhibitors (TIMPs), causing LVH and therefore heart failure and death [71,72].

Searched in the literature for evidence of a causal association between low vitamin D levels and increased risk of cardiovascular disease, based on Hill criteria and found that all relevant Hill criteria for a causal association in a biological system have been met, to indicate that vitamin deficiency is a risk factor for cardiovascular disease [72].

The strength of this study was the reduced loss of data due to the lack of records in the medical records, as well as the use of a wide range of complementary tests, investigating several aspects of the health conditions of the elderly, as well as making it possible to evaluate the cardiological aging to guide the conduct therapy.

All patients in this study underwent Doppler echocardiography and the findings of left ventricular hypertrophy and reduction of LV function analysed by this method, were associated with vitamin D deficiency, demonstrating the importance of this test, due to alterations in the structure and cardiac function in the elderly are usually subclinical, which often precede the development of heart failure.

Based on the evidence presented in the present study, supported by the literature, it can be stated that the high percentage of elderly people with vitamin D deficiency and their association with cardiovascular risk factors suggests that vitamin D recommendations should be implemented especially in services of basic health care. The ease of quantification of vitamin D, the low cost of its supplementation and the possibility of preventing and treating cardiovascular diseases seem to point to the need for a greater amount of studies of vitamin D supplementation in a prospective cohort, so that the supplementation conduct is implanted with a solid evidence base.

Among the limitations of this study is the transversal nature, which makes it impossible to establish causal relationships, as well as the fact that it was performed through medical records analyzes, since it did not contain

more detailed information such as physical exercise practice, quantification of patients who used of vitamin D supplementation. However, this limitation was reduced because the data collection was done in specialized places to care for the elderly, maintaining a broad and systematized research routine, with better cardiovascular risk assessment conditions. It was also not possible to identify bone densitometry, serum calcium, phosphorus and PTH dosages in all patient charts of this study, and the presence of osteoporosis in the elderly could not be quantified, and well better analysis of bone metabolism

Another limitation of the present study was the no quantification of LV diastolic dysfunction in the elderly, since it is a very frequent alteration in this age range [73,74].

Conclusions

A high prevalence of vitamin D deficiency was detected in this population of elderly people. These results were significantly associated in this study with males and the age range ≤ 70 years, in the same way associated with cardiovascular risk factors such as smoking, obesity, DMII, dementia, depression, cardiac arrhythmia and, mainly, a strong association with CAD and hypertrophy or reduction of left ventricular function.

Observational data support the relationship between vitamin D levels and cardiovascular disease and therefore vitamin D deficiency can be considered a marker of cardiovascular risk. Large randomized controlled trials are needed to establish protocols to establish which optimal levels of 25-Hydroxyvitamin D are recommended to maintain a healthy profile in patients with CVD. This maintenance of an ideal serum vitamin D level is of fundamental importance, not only for regulation of calcium balance, but also to decrease cardiovascular risk, to control systemic arterial pressure, inhibiting the progression of atherosclerosis and consequently preventing CAD and peripheral arterial disease, as a neuroprotective agent in the prevention of stroke, dementia and depression and controlling the constituent elements of the metabolic syndrome.

The evidence presented in this study, supported by the literature, may be relevant for primary health care, considering the high prevalence of vitamin D deficiency in the elderly population, as well as being a treatable condition with low cost supplementation. Vitamin D supplementation may be combined with antihypertensives or antidiabetics as a simple prophylactic agent for cardiovascular morbidities, especially in the elderly, where small gains in prevention are important from a public health point of view.

These actions can be very important in guiding public health policies, considering the high prevalence of vitamin D deficiency and aging of the population. The benefits of this public health approach to vitamin D assessment and treatment may probably reflect a reduction in cardiovascular morbidity and mortality.

Declaration of Interest

The authors report no conflicts of interest. The authors alone are responsible for writing the paper and the contents therein.

References

- Gravina CF, Franken R, Wenger N, Freitas EV, Batlouni M, et al. (2010) II diretrizes em cardiogeriatría da Sociedade Brasileira de Cardiologia. *Arq Bras Cardiol* 95: e16-e76.
- Sociedade Brasileira de Nefrologia (2010) VI Brazilian Guidelines on Hypertension. *Arq Bras Cardiol* 95: 1-51.
- Giovannucci E, Liu Y, Hollis BW, Rimm EB (2008) 25-hydroxyvitamin D and risk of myocardial infarction in men: a prospective study. *Arch Intern Med* 168: 1174-1180.
- Anderson JL, May HT, Horne BD, Bair TL, Hall NL, et al. (2010) Relation of vitamin D deficiency to cardiovascular risk factors, disease status, and incident events in a general healthcare population. *Am J Cardiol* 106: 963-968.
- Karakas M, Thorand B, Zierer A, Huth C, Meisinger C, et al. (2013) Low levels of serum 25-Hydroxyvitamin D are associated with increased risk of myocardial infarction, especially in women: results from the MONICA/KORA Augsburg case-cohort study. *J Clin Endocrinol Metab* 98: 272-280.
- Tomson J, Emberson J, Hill M, Gordon A, Armitage J, et al. (2013) Vitamin D and risk of death from vascular and non-vascular causes in the whitehall study and meta-analyses of 12.000 death. *Eur Heart J* 34: 1365-1374.
- Oz F, Cizgici AY, Oflaz H, Elitok A, Karaayvaz EB, et al. (2013) Impact of vitamin D insufficiency on the epicardial coronary flow velocity and endothelial function. *Coron Artery Dis* 24: 392-397.
- Ginde AA, Scragg R, Schwartz RS, Camargo CA Jr (2009) Prospective study of serum 25-hydroxyvitamin D level, cardiovascular disease mortality, and all-cause mortality in older U.S. adults. *J Am Geriatr Soc* 57: 1595-1603.
- Perna L, Schöttker B, Holleczer B, Brenner H (2013) Serum 25-hydroxyvitamin D and incidence of fatal and non-fatal cardiovascular events: a prospective study with repeated measurements. *J Clin Endocrinol Metab* 98: 4908-4915.
- Theodoratou E, Tzoulaki I, Zgaga L, Ioannidis JP (2014) Vitamin D and multiple health outcomes: umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ* 348: g2035.
- Hosseini-Nezhad A, Michael F, Holick MF (2013) Vitamin D for Health: a global perspective. *Mayo Clin Proc* 88: 720-755.
- Henry HL, Bouillon R, Norman AW, Gallagher JC, Lips P, et al. (2010) 14th Vitamin D Workshop consensus on vitamin D nutritional guidelines. *J Steroid Biochem Mol Biol* 121: 4-6.
- Khaw KT, Luben R, Wareham N (2014) Serum 25-hydroxyvitamin D, mortality, and incident cardiovascular disease, respiratory disease, cancers, and fractures; a 13-y prospective populations study. *Am J Clin Nutr* 100: 1361-7130.
- Pareja-Galeano H, Alis R, Sanchis-Gomar F, Lucia A, Emanuele E (2015) Vitamin D, precocious acute myocardial infarction, and exceptional longevity. *Int J Cardiol* 199: 405-406.
- Franco I (1994) It deals with the national policy of the elderly, creates the National Council for the Elderly and provides other measures, [http://www.planalto.gov.br/ccivil_03/leis/L8842.htm]. 20 September 2016.
- Hildrum B, Mykletun A, Hole T, Midthjell K, Dahl AA (2007) Age-specific prevalence of the metabolic syndrome defined by the International Diabetes Federation and the National Cholesterol Education Program: the Norwegian HUNT 2 study. *BMC Public Health* 7: 220.
- Maeda SS, Saraiva GL, Kunni IS, Hayashi LF, Cendoroglo MS, et al. (2013) Factors affecting vitamin D status in different populations in the city of São Paulo, Brazil: the São Paulo vitamin D valuation Study (SPADES). *BMC Endocr Disord* 13: 14.
- Martins D, Wolf M, Pan D, Zadshir A, Tareen N, et al. (2007) Prevalence of cardiovascular risk factors and the serum levels of 25-hydroxyvitamin D in the United States: data from the third National Health and Nutrition Examination Survey. *Arch Int Med* 167: 1159-1165.

19. Souberbielle JC, Massart C, Brailly-Tabard S, Cavalier E, Chanson P (2016) Prevalence and determinants of vitamin D deficiency in healthy French adults: the VARIETE study. *Endocrine* 53: 543-550.
20. Raina AH, Allai MS, Shah ZA, Changal KH, Raina MA, et al. (2016) Association of low levels of vitamin D with chronic stable angina: a prospective case-control study. *N Am J Med Sci* 8: 143-150.
21. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, et al. (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an endocrine society clinical practice guideline. *J Clin Endocr Metabol* 96: 1911-1930.
22. American Geriatrics Society Consensus Statement (2014) Vitamin D for prevention of falls and their consequences in older Adults year of publication. American Geriatrics Society.
23. Lavie CJ, Lee JH, Milani RV (2011) Vitamin D and cardiovascular disease will it live up to its hype? *J Am Coll Cardiol* 58: 1547-1556.
24. Chaimowicz F and Camargos MCS (2011) Envelhecimento e saúde no Brasil. In: Freitas EV, Py L, editores. *Tratado de geriatria e gerontologia*. 3. ed. Rio de Janeiro: Guanabara Koogan, 74-98.
25. Penna G and Adorini L (2000) $1\alpha,25$ -Dihydroxyvitamin D₃ Inhibits Differentiation, Maturation, Activation, and Survival of Dendritic Cells Leading to Impaired Alloreactive T Cell Activation. *J Immunol* 164: 2405-2411.
26. Holick MF (2007) Vitamin D deficiency. *N Engl J Med* 357: 266-281.
27. Vogt S, Baumert J, Peters A, Thorand B, Scragg R (2016) Waist circumference modifies the association between serum 25(OH)D and systolic blood pressure: results from NHANES 2001-2006. *J Hypertens* 34: 637-645.
28. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF (2000) Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 72: 690-693.
29. Pittas AG, Harris SS, Stark PC, Dawson-Hughes B (2007) The effects of calcium and vitamin D supplementation on blood glucose and markers of inflammation in nondiabetic adults. *Diabetes Care* 30: 980-986.
30. Gradinaru D, Borsa C, Ionescu C, Margina D, Prada GI, et al. (2012) Vitamin D status and oxidative stress markers in the elderly with impaired fasting glucose and type 2 diabetes mellitus. *Aging Clin Exp Res* 24: 595-602.
31. Kuloğlu O, Gür M, Şeker T, Kalkan GY, Kırım S, et al. (2013) Serum hydroxyvitamin D level is associated with aortic distensibility and left ventricle hypertrophy in newly diagnosed type 2 diabetes mellitus. *Diab Vasc Dis Res* 10: 546-549.
32. Raska J Jr, Rašková M, Zikán V, Škrha J (2016) High prevalence of hypovitaminosis D in postmenopausal women with type 2 diabetes mellitus. *Prague Med Rep* 117: 5-17.
33. Forouhi NG, Ye Z, Rickard AP, Khaw KT, Luben R et al. (2012) Circulating 25-hydroxyvitamin D concentration and the risk of type 2 diabetes: results from the European Prospective Investigation into Cancer (EPIC)-Norfolk cohort and updated meta-analysis of prospective studies. *Diabetologia* 55: 2173-2182.
34. Llewellyn DJ, Lang IA, Langa KM, Melzer D (2011) Vitamin D and cognitive impairment in the elderly US population. *J Gerontol A Biol Sci Med Sci* 66: 59-65.
35. Przybelski RJ and Binkley NC (2007) Is vitamin D important for preserving cognition? A positive correlation of serum 25-hydroxyvitamin D concentration with cognitive function. *Arch Biochem Biophys* 460: 202-205.
36. Annweiler C, Fantino B, Schott AM, Krolak-Salmon P, Allali G, et al. (2012) Vitamin D insufficiency and mild cognitive impairment: cross-sectional association. *Eur J Neurol* 19: 1023-1029.
37. Balion C, Griffith LE, Striffler L, Henderson M, Patterson C, et al. (2012) Vitamin D, cognition, and dementia: a systematic review and meta-analysis. *Neurology* 79: 1397-1405.
38. Keeny JT and Butterfield DA (2015) Vitamin D deficiency and Alzheimer disease: common links. *Neurobiol Dis* 84: 84-98.
39. Lett HS, Blumenthal JA, Babyak MA, Sherwood A, Strauman T, et al. (2004) Depression as a risk factor for coronary artery disease: evidence, mechanism, and treatment. *Psychosom Med* 66: 305-315.
40. Wilkins CH, Sheline YI, Roe CM, Birge SJ, Morris JC (2006) Vitamin D deficiency is associated with low mood and worse cognitive performance in older adults. *Am J Psychiatry* 14: 1032-1040.
41. Armstrong DJ, Meenagh GK, Bickle I, Lee AS, Curran ES, et al. (2007) Vitamin D deficiency is associated with anxiety and depression in fibromyalgia. *Clin Rheumatol*, 26: 551-554.
42. Jorde R, Waterloo K, Saleh F, Haug E, Svartberg J (2006) Neuropsychological function in relation to serum parathyroid hormone and serum 25-hydroxyvitamin D levels. *J. Neurol* 253: 464-470.
43. Hoogendijk WJ, Lips P, Dik MG, Deeg DJ, Beekman AT, et al. (2008) Depression is associated with decreased 25-hydroxyvitamin D and increased parathyroid hormone levels in older adults. *Arch Gen Psychiatry* 65: 508-512.
44. Yeshokumar AK, Saylor D, Kornberg MD, Mowry EM (2015) Evidence for the importance of vitamin D status in neurologic conditions. *Curr Treat Options Neurol* 17: 51.
45. Spedding S (2014) Vitamin D and depression: A systematic review and meta-analysis comparing studies with and without biological flaws. *Nutrients* 6: 1501-1518.
46. Moy FM, Hoe VC, Hairi NN, Vethakkan SR, Bulgiba A (2017) Vitamin D deficiency and depression among women from an urban community in a tropical country. *Public Health Nutr* 20: 1844-1850.
47. Frost L, Søren, PJ, Pedersen L, Husted S, Engholm G, et al. (2002) Seasonal variation in hospital discharge diagnosis in atrial fibrillation: a population-based study. *Epidemiology* 13: 211-215.
48. Murphy NF, Stewart S, MacIntyre K, Capewell S, McMurray JJ (2004) Seasonal variation in morbidity and mortality related to atrial fibrillation. *Int J Cardiol* 97: 283-288.
49. Demir M, Uyan U, Melek M (2014) The effects of vitamin D deficiency on atrial fibrillation. *Clin Appl Thromb Hemost* 20: 98-103.
50. Chen WR, Liu ZY, Shi Y, Yin DW, Wang H, et al. (2014) Relation of low vitamin D to nonvalvular persistent atrial fibrillation in chinese patients. *Ann Noninvasive Electrocardiol* 19: 166-173.
51. Lappegard KT, Hovland A, Pop GA, Mollnes TE (2013) Atrial fibrillation: inflammation in disguise? *Scand J Immunol* 78: 112-119.
52. Hafany DA, Chang S, Lu Y, Chen Y, Kao Y, et al. (2014) Electromechanical effects of 1,25-dihydroxyvitamin D with antiatrial fibrillation activities. *J Cardiovasc Electrophysiol* 25: 317-323.
53. Rienstra M, Cheng S, Larson MG, McCabe EL, Booth SL, et al. (2011) Vitamin D status is not related to development of atrial fibrillation in the community. *Am Heart J* 62: 538-541.
54. Qayyum F, Landex NL, Agner BR, Rasmussen M, Jøns C, et al. (2012) Vitamin D deficiency is unrelated to type of atrial fibrillation and its complications. *Dan Med J* 59: A4505.
55. Vitezova A, Cartolano NS, Heeringa J, Zillikens MC, Hofman A, et al. (2015) Vitamin D status and atrial fibrillation in the elderly: the rotterdam study. *PLoSOne* 10: e0125161.
56. Kunadian V, Ford GA, Bawamia B, Qiu W, Manson JE (2014) Vitamin D deficiency and coronary artery disease: a review of the evidence. *Am Heart J* 167: 283-291.
57. Ipek G, Kurmus O, Koseoglu C, Onuk T, Gungor B, et al. (2017) Predictors of in-hospital mortality in octogenarian patients who underwent primary percutaneous coronary intervention after ST segment elevated myocardial infarction. *Geriatr Gerontol Int* 17: 584-590.
58. Wang TJ, Pencina MJ, Booth SL, Jacques PF, Ingelsson E, et al. (2008) Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 117: 503-511.

59. Kestenbaum B, Katz R, de Boer I, Hoofnagle A, Sarnak MJ, et al. (2011) Vitamin D, parathyroid hormone, and cardiovascular events among older adults. *J Am Coll Cardiol* 58: 433-441.
60. Cure MC, Cure E, Yuce S, Yazici T, Karakoyun I, et al. (2014) Mean platelet volume and vitamin D level. *Ann Lab Med* 34: 98-103.
61. Fall T, Shiue I, Bergeå af Geijerstam P, Sundström J, Årnlöv J, et al. (2012) Relations of circulating vitamin D concentrations with left ventricular geometry and function. *Eur J Heart Fail* 14: 985-991.
62. Li YC (2003) Vitamin D regulation of the renin-angiotensin system. *J Cell Biochem* 88: 327-331.
63. Xiang W, Kong J, Chen S, Cao LP, Qiao G, et al. (2005) Cardiac hypertrophy in vitamin D receptor knockout mice: role of the systemic and cardiac renin-angiotensin systems. *Am J Physiol Endocrinol Metab* 288: E125-E132.
64. Mancuso P, Rahman A, Hershey SD, Dandu L, Nibbelink KA, et al. (2008) 1,25-Dihydroxyvitamin D3 treatment reduces cardiac hypertrophy and left ventricular diameter in spontaneously hypertensive heart failure prone(cp/+) rats independent of changes in serum leptin. *J Cardiovasc Pharmacol* 51: 559-564.
65. Zittermann A and Koefer R (2008) Vitamin D in the prevention and treatment of coronary heart disease. *Curr Opin Clin Nutr Metab Care* 11: 752-757.
66. Helming L, Böse J, Ehrchen J, Schiebe S, Frahm T, et al. (2005) 1 α ,25-dihydroxyvitamin D3 is a potent suppressor of interferon gamma-mediated macrophage activation. *Blood* 106:4351-4358.
67. Rahman A, Hershey S, Ahmed S, Nibbelink K, Simpson RU (2007) Heart extracellular matrix gene expression profile in the vitamin D receptor knockout mice. *J Steroid Biochem Mol Biol* 103: 416-419.
68. Santoro D, Gagliostro G, Alibrandi A, Ientile R, Bellinghieri G, et al. (2014) Vitamin D receptor gene polymorphism and left ventricular hypertrophy in chronic kidney disease. *Nutrients* 10: 1029-1037.
69. Porto CM, Silva VDL, da Luz JSB, Filho BM, da Silveira VM (2017) Association between vitamin D deficiency and heart failure risk in the elderly. *ESC Heart Fail* 5: 63-74.
70. Liu M, Li X, Sun R, Zeng YI, Chen S, et al. (2016) Vitamin D nutritional status and the risk for cardiovascular disease. *Exp Ther Med* 11: 1189-1193.
71. Polyakova V, Loeffler I, Hein S, Miyagawa S, Piotrowska I, et al. (2011) Fibrosis in endstage human heart failure: severe changes in collagens metabolism and MMP/TIMP profiles. *Int J Cardiol* 151: 18-33.
72. Weyland PG, Grant WB, Howie-Esquivel J (2014) Does sufficient evidence exist to support a causal association between vitamin D status and cardiovascular disease risk? An Assessment Using Hill's Criteria for Causality. *Nutrients* 6: 3403-3430.
73. Moutinho MAE, Colucci FA, Alcoforado V, Tavares LR, Rachid MBF, et al. (2008) Heart failure with preserved ejection fraction and systolic dysfunction in the community. *Arq Bras Cardiol* 90: 132-137.
74. Goldraich L, Clausell N, Biolo A, Beck-da-Silva L, Rohde LE, et al. (2010) Preditores clínicos de fração de ejeção de ventrículo esquerdo preservada na insuficiência cardíaca descompensada. *Arq Bras Cardiol* 94: 364-371.